

*In The Name of God*



Ardabil University of  
Medical Sciences

**School of Medicine**

Submitted to the School of Medicine in Partial Fulfillment of the Requirements  
for the Degree of Master of Science (M.Sc.) in Clinical Biochemistry

Subject:

**The Effect of Dimethyl Sulfone on Gastrointestinal Cancer  
Cell Lines (AGS, HepG2 and KYSE-30) and Flow Cytometric  
Analysis of the Cell Cycle**

Supervisor:

**Dr. Shahab Bohlooli**

Advisor:

**Dr. Mohammad Mazani**

**By: Naser Jafari**

## Dedication

Dedicated to:

My Unfailing Kind *Mother*

Who has been supportive of me in every aspect of life,

And to My Brother and Sister:

*Saeed* and *Mahdieh*

For their kindness

And

To my kind professor *Dr. Bohlooli*,

A brilliant researcher, scientist, and teacher,  
For training me to become a researcher, I have learned  
so much in the lab, it is impossible to recount the  
details in this space.

Thanks for all of your help and advice, and for  
providing me with so many opportunities to discover.

To researchers and scientists, who with their hard work  
pave the road to discovery,

Last but not least, to anyone who has been touched by  
cancer, as they are my motivation for this thesis and for  
my research.

## **Acknowledgments**

My sincere thanks are due to the friends and people, who made my research possible,

Including gentlemen: Khanzadeh, Hosseini, Rostami, Mohammadnia, Khodayi, Khaligh, Shahbazzadegan,

And ladies: Naftchi, Golmaghani, Esmaeeli, Molazadeh and Mohandesi

Thank you all

I am grateful of my friends, who made my life outside of lab exciting,

Saeed Aslani, Poorya Badihi, Aboozar Barzegar, Rahim Delsouzi, Nemat Heydarogloo, Naser Jafargholizadeh, Yaser Khakpour, Amir Mahmoodzadeh, Yashar Nabizadeh, Mehrdad Nickmanesh, Rezgar Rahbari, Mehrdad Sarvghamat, Ali Shadmanfar, Hasan Shaghaghi, Saeed Zarenejad.

Thanks for everything,

## **Abstract:**

**Background:** Methylsulfonylmethane (MSM) is a dietary supplement to support healthy body. It is considered to be one of the least toxic substances in biology. Considering preventive effects of this substance on tumor onset and its low toxicity to healthy body we investigated in vitro cytotoxic effects of methylsulfonylmethane on cancer cell lines. **Methods:** AGS, HepG2 and KYSE-30 cancer cell lines were treated by MSM by initial concentration of 50 mg/ml with  $\frac{1}{4}$  serial dilutions and incubated for 24, 48 and 72 hours. Cytotoxicity was examined through MTT, neutral red uptake and protein measurement assays. EB/AO staining was used for apoptotic cell detection. A DAPI staining method was used to analysis cell cycle by flow cytometry. **Results:** IC<sub>50</sub> of the MSM on AGS, HepG2 and KYSE-30 cell lines were 28.04, 21.87 and 27.98 mg/ml after 72 hours respectively. The EB/AO staining showed an increase in apoptotic cells after MSM treatment. The flow cytometry cell cycle analysis showed a significant increase in cell density at G2/M phase. **Conclusion:** MSM had cytotoxic effect on cancer cell lines but HepG2 cell line was more susceptible. This study provides the evidence that MSM may induce cytotoxic effect on cancer cell lines by apoptosis and it could be of value in realm of chemotherapy.

**Key Words:** Methylsulfonylmethane, MSM, Cancer cell lines, Cytotoxicity, Apoptosis, Adjuvant

## Abbreviations:

Abbreviation	Text
°C	Centigrade
µg	Micro-gram
µl	Micro-liter
AO	Acridine Orange
BBB	Blood Brain Barrier
CAT	Catalase
cm	Centimeter
COX	Cyclooxygenase
DAPI	4',6'-Diamidino-2-Phenylindole
DMSO	Dimethyl Sulfoxide
DMSO <sub>2</sub>	Dimethyl Sulfone
DNA	Deoxyribonucleic Acid
EB	Ethidium Bromide
FBS	Fetal Bovine Serum
FDA	Food and Drug Administration
FRAME	Fund for the Replacement of Animals in Medical Experiments

g	Gram
G1	Gap 1 Phase
G2	Gap 2 Phase
GC	Gastric Cancer
GI	Gastrointestinal
GSH	Glutathione
h	Hour(s)
HBV	Hepatitis B Virus
HCC	Hepatocellular Carcinoma
HCV	Hepatitis C Virus
IC	Interstitial Cystitis
IC50	The Half Maximal Inhibitory Concentration
Ig E	Immunoglobulin E
IL-1 $\beta$	Interleukin 1 Beta
IV	Intravenous
kPa	Kilopascal
kg	Kilogram
LDL	Low Density Lipoprotein
M	Mitosis



MDA	Malondialdehyde
mg	Milligram
min	Minute
ml	Milliliter
mol wt	Molecular Weight
MPO	Myeloperoxidase
MSM	Methylsulfonylmethane
MTT	MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5 Diphenyltetrazolium Bromide)
NAD(P)H	Nicotinamide Adenine Dinucleotide (Phosphate)
NIH	National Institute of Health
nm	Nanometer
OD	Optical Density
PBS	Phosphate Buffered Saline
pg	Pico-gram
PGE2	Prostaglandin E2
ppm	Parts Per Million
rcf	Relative Centrifugal Force
rpm	Round Per Minute
S	Synthesis Phase

SD	Standard Deviation
SEM	Standard Error of the Mean
UV	Ultraviolet
w/v	Weight/Volume

## Figures and Tables

Figure 1- 1 Overview of chnges in cells that cause cancer .....	2
Figure 1- 2 Gastrointestinal system .....	5
Figure 1- 3 The fate of a single parental chromosome throughout the eukaryotic cell-cycle.....	11
Figure 1- 4 Mitosis and cell-cycle transition.....	13
Figure 1- 5 Progression of cells through the various phases of mitosis .....	15
Figure 1- 6 Cell-cycle checkpoints .....	20
Figure 1- 7 Apoptotic cells. Apoptotic cells break apart without spewing forth cell constituents that might harm neighboring cells.....	22
Figure 1- 8 Dimethyl sulfone.....	25
Figure 3-1 Major equipment.....	39
Figure 3- 2 AGS cell line.....	41
Figure 3- 3 HepG2 cell line .....	42
Figure 3- 4 KYSE-30 cell line .....	43
Figure 3- 5 The hemocytometer.....	46
Figure 4- 1 MTT assay of cytotoxic activity of MSM on AGS, HepG2 and KYSE-30 cell lines after 24 hours compared to normal .....	58
Figure 4- 2 MTT assay of cytotoxic activity of MSM on AGS, HepG2 and KYSE-30 cell lines after 48 hours compared to normal .....	59
Figure 4- 3 MTT assay of cytotoxic activity of MSM on AGS, HepG2 and KYSE-30 cell lines after 72 hours compared to normal .....	60
Figure 4- 4 Neutral Red (NR) assay of cytotoxic activity of MSM on AGS, HepG2 and KYSE-30 cell lines after 24 hours compared to normal .....	61
Figure 4- 5 Neutral Red (NR) assay of cytotoxic activity of MSM on AGS, HepG2 and KYSE-30 cell lines after 48 hours compared to normal .....	62
Figure 4- 6 Neutral Red (NR) assay of cytotoxic activity of MSM on AGS, HepG2 and KYSE-30 cell lines after 72 hours compared to normal .....	63

Figure 4- 7 Protein Measurement (FRAME) assay of cytotoxic activity of MSM on AGS, HepG2 and KYSE-30 cell lines after 24 hours compared to normal .....	64
Figure 4- 8 Protein Measurement (FRAME) assay of cytotoxic activity of MSM on AGS, HepG2 and KYSE-30 cell lines after 48 hours compared to normal .....	65
Figure 4- 9 Protein Measurement (FRAME) assay of cytotoxic activity of MSM on AGS, HepG2 and KYSE-30 cell lines after 72 hours compared to normal .....	66
Figure 4- 10 Apoptosis induction of MSM on AGS cell line .....	69
Figure 4- 11 The histogram of percentage of cell death in AGS cells after treating with MSM .....	70
Figure 4- 12 Apoptosis induction of MSM on HepG2 cell line .....	71
Figure 4- 13 The histogram of percentage of cell death in HepG2 cells after treating with MSM .....	72
Figure 4- 14 Apoptosis induction of MSM on KYSE-30 cell line.....	73
Figure 4- 15 The histogram of percentage of cell death in KYSE-30 cells after treating with MSM .....	74
Figure 4- 16 Effect of MSM on cell-cycle distribution in AGS cell line .....	77
Figure 4- 17 Effect of MSM on cell-cycle distribution in HepG2 cell line.....	78
Figure 4- 18 Effect of MSM on cell-cycle distribution in KYSE-30 cell line...	79
 Table 4- 1 IC50 values of MSM on AGS, KYSE-30 and HepG2 cell lines after 24, 48 and 72 h determined by MTT, neutral red and frame assays .....	57
Table 4- 2 Apoptosis induction of MSM on AGS, KYSE-30 and HepG2 cell lines data are expressed as percentage of cells .....	68
Table 4- 3 Effect of MSM on cell-cycle distribution on AGS, KYSE-30 and HepG2 cell lines .....	76

## Table of Contents

Dedication .....	c
Acknowledgments .....	e
Abstract: .....	f
Abbreviations: .....	g
Figures and Tables .....	k
1: Introduction .....	1
1.1 Cancer: .....	1
1.1.1 Gastrointestinal Cancers .....	3
1.2 Cell-Cycle .....	10
1.3 Mitosis .....	14
1.3.1 Interphase and Mitosis .....	14
1.3.2 Mitotic Stages .....	16
1.4 Cell-Cycle Checkpoints .....	18
1.5 The Cell-Cycle and Cancer .....	20
1.6 Cell Death .....	21
1.6.1 Apoptosis .....	23
1.6.2 Necrosis .....	24
1.7 Dimethyl Sulfone .....	24

1.7.1 The Discovery of MSM: .....	26
1.7.2 Biochemistry of Dimethyl Sulfone .....	27
1.8 Objectives and Hypotheses .....	28
1.8.1 Objectives and Hypotheses .....	28
1.8.2 Special objectives .....	28
1.8.3 Hypotheses .....	28
2. Literature Review .....	29
2.1 Mechanism of Action of MSM: .....	29
2.2 Clinical Indications.....	30
2.2.1 Interstitial Cystitis: .....	30
2.2.2 Ulcerative Colitis.....	31
2.2.3 Allergic Rhinitis: .....	31
2.2.4 Chemopreventive Properties:.....	32
2.2.5 Autoimmune Disease:.....	33
2.2.6 Other Clinical Indications:.....	33
2.3 Side Effects and Toxicity: .....	34
2.4 Dosage: .....	34
2.5 Study Design: .....	34
3. Materials and Methods .....	35
3.1 Materials and Equipment:.....	35
3.1.1 Reagents: .....	35

3.1.2 Dishes and Disposables: .....	36
3.1.3 Equipment: .....	37
3.2 Methods: .....	40
3.2.1 Cell culture and treatment:.....	40
3.2.2 Viability Measurements.....	48
3.2.3 EB/AO staining for apoptotic cells: .....	52
3.2.4 Cell-Cycle Analysis by Flow Cytometry: .....	53
3.3 Statistical analysis: .....	55
4. Results:.....	56
4.1. Growth inhibition study:.....	56
4.2. EB/AO staining for apoptotic cells: .....	67
4.3. Cell-Cycle Analysis by Flow Cytometry: .....	75
5. Discussion: .....	80
5.1 Conclusion: .....	84
5.2 Limitations: .....	85
5.3 Suggestions: .....	86
6. References:.....	87

# **1: Introduction**

## **1.1 Cancer:**

Cancer causes about one-fifth of the deaths each year. About 100-350 of 100,000 people scarify by cancer in a year. Cancer is due to failure of the mechanisms that usually control the growth and proliferation of cells. Throughout adult life and during normal development, genetic control systems regulate the balance between cell birth and death in response to various signals. Cell birth and death rates determine adult body size. The cells in many adult tissues normally do not proliferate except during healing processes. Such stable cells (e.g., hepatocytes, gastric and intestinal epithelial cells, and white blood cells) can remain functional for long periods or even for the entire lifetime of an organism. Cancer occurs when the mechanisms that maintain these normal growth rates malfunction to cause excess cell division. The failure of cellular regulation that give rise to most or all cases of cancer are due to genetic damage that is often accompanied by influences of tumor-promoting chemicals, hormones and sometimes viruses [1] ( Figure 1-1).